

OBESITY AND ASTHMA: NUTRITION RISK FACTORS IN ADOLESCENTS

Franceliane Jobim Benedetti^{1*}, Vera Lúcia Bosa², Juliana Mariante Giesta³, Gilberto Bueno Fischer⁴

1. Nutritionist, Master's Graduate Program in Health child and adolescent in Universidade Federal do Rio Grande do Sul (UFRGS) and Professor Graduate in Centro Univeritário Franciscano.
2. Nutritionist, Ph.D. in Health of Children and Adolescents, Professor Graduate in Universidade Federal do Rio Grande do Sul (UFRGS).
3. Nutritionist, Graduate Program of Multiprofessional Residency in Health, concentration in Child and Adolescent Health of the Hospital de Clínicas de Porto Alegre (HCPA).
4. Ph.D. in Medicine Pneumology, Professor of Pediatrics at the Universidade Federal de Ciências da Saúde de Porto Alegre (UFCSPA) and Professor Graduate Program in Medical Science-Pediatrics (UFRGS).

ABSTRACT:

The objective of this study was to confirm the nutritional risk factors for asthma and obesity in adolescents using a cross-sectional survey. We included male and female adolescents aged 10 to 19 years of age in this study. The indicators of nutritional status used in this study were body mass index, the height-for-age index, and arm and waist circumferences. We used a food frequency questionnaire to estimate the average consumption of sodium. Physical activity was assessed using the International Physical Activity Questionnaire, and asthma characteristics were evaluated using the International Study of Asthma and Allergies in Childhood (ISAAC). Factors associated with asthma in bivariate and multivariate analyses were assessed using the Poisson regression analysis. The significance level was set at 5%, and the analyses were performed using the SPSS statistical package. We evaluated 1362 students with an average age of 15.65 ± 1.24 years. There was a positive risk for asthma between females (PR = 1.41) and asthma severity in adolescents with a high sodium intake (PR = 2.30). Within overweight adolescents, the prevalence for asthma risk was higher among females (PR = 1.66) and the high sodium intake group (PR = 1.98). An increased risk for asthma severity correlated with high sodium intake (PR = 3.07). There was a higher risk for asthma and its severity in females with high sodium intake. The same pattern was observed for adolescents with excessive weight.

Corresponding author: Franceliane Jobim Benedetti – Street Ramiro Barcelos, 2400 - 2nd floor, District Santana, CEP 90035-00. Porto Alegre, RS-Brazil. Phone: (55) 33476496 Fax: (51) 32148646 E-mail address: francijb@yahoo.com.br

Citation: Franceliane Jobim Benedetti, Vera Lúcia Bosa, Juliana Mariante Giesta, Gilberto Bueno Fischer (2016) Obesity and Asthma: Nutrition Risk Factors In Adolescents. International Journal of Nutrition - 1(3):47-60. <https://doi.org/10.14302/issn.2379-7835.ijn-15-770>

Keywords: adolescents; asthma; obesity; sodium; physical activity.

Received 05 Oct 2015;

Accepted 20 Dec 2015;

Published 11 Jan 2016

Academic Editor: Justin Friedlander, Fox Chase/Einstein Urologic Institute

INTRODUCTION

Asthma and obesity are diseases that have reached epidemic proportions, especially in westernized countries. Theories have been proposed to explain the association between obesity and asthma. Theoretical assumptions included the direct effects of obesity on functional, genetic, hormonal, and immunological mechanisms and inflammatory responses (1-3). Moreover, increased weight has been associated with lifestyle and eating habits (4, 5).

Significant changes in dietary and nutritional standards have been proposed worldwide. These changes are characterized as part of a process called nutritional transition. There is an increasing consumption of processed foods and a more sedentary lifestyle provided by technological advances, especially among children and adolescents who remain in front of televisions, videogames and computer devices for many hours (6, 7). The low level of physical activity in adolescents with asthma may be associated with obesity because the severity of the disease can contribute to a sedentary lifestyle; however, obesity and a large amount of time spent watching TV can increase the risk of respiratory symptoms (8, 9).

In addition to the relationship with respiratory health and sedentary behavior, the habit of watching television promotes a change in eating routine and increases the risk of obesity in children and adolescents (10). These habits are also associated with a higher consumption of salty snacks (11) that are rich in sodium and increase the risk of bronchial hyperresponsiveness and wheezing in children and adolescents (12).

Despite the promising discoveries and hypotheses available in the current literature, finding on the prevalence of asthma, the effects of a sedentary lifestyle and the role of sodium are scarce.

Therefore, this study assessed the nutritional risk factors for asthma and obesity in adolescents.

METHODOLOGY

This study consisted of cross-sectional observational design and was completed with a group of adolescents from 16 public high schools located in the urban area of the city of Santa Maria-RS/Brazil from May to November 2012.

We included male and female adolescents aged 10 and 19 years who were regularly enrolled in the schools where data collection occurred. All subjects agreed to participate, and their parents signed an informed consent form. We excluded three adolescents with cerebral palsy and two adolescents with Down syndrome.

Sampling was performed by randomly polling classes. The sample size estimation to show statistically significant differences with a RR= 2.0 for the prevalence of obesity (15%) among adolescents with and without asthma (21%), a significance level of 0.05 and a power level of 80% was calculated as 395 individuals.

The research project used to develop this study was approved by the Ethics Committee in research of the Rio Grande do Sul Federal University (UFRGS) under the protocol n. 20009. All participants received the results of their assessment, and the school received the overall results of the student analyses.

The data referring to sample characterizations, anthropometric evaluations and asthma characteristics were collected by the authors and previously trained students from the undergraduate program in nutrition. The questionnaires were applied in the classroom for all of the present students and in the absence of the teacher.

Anthropometric measurements were assessed individually and in duplicate using standard techniques

and calibrated equipment in a private room. We assumed a maximum difference value of 1.0cm or 100g between measurements. For the analysis, we used the average value of each measure (13).

The body mass weight in kg was obtained while the adolescents were barefoot and wearing minimal clothing using platform scales (Plenna, São Paulo, Brazil) with a maximum load of 150kg and a range of 100g. Height was measured while the subjects were standing barefoot using an extensible stadiometer (Sanny, São Paulo, Brazil) fixed on a flat wall without a baseboard (13).

The anthropometric indicators used included the body mass index for age (BMI-Z) and the height-for-age index (H/A-Z). These parameters were analyzed with *Anthro Plus* version 1.04, and the results were obtained using the z-score and classified according to WHO guidelines (13). Adolescents with a BMI-Z greater than +1 were classified as overweight.

For arm circumference (AC) measurements, we used an inextensible measuring tape (Secca, ON, USA) that was positioned perpendicular to the long axis of the arm at the midpoint (distance between the acromion and the olecranon). The measurement value was classified based on Frisancho (14).

Waist circumference (WC) was measured using a Secca® (ON, USA) in elastic measuring tape positioned at the minimum circumference between the iliac crest and the last rib. An excess of abdominal fat was defined as a WC higher than the 80th percentile for age and sex (15).

We used a food frequency questionnaire for foods with high sodium content (FFQSo) to estimate the average daily intake of sodium. The survey inquires about 15 types of food, including the frequency and portion size consumed. The final result was calculated as the average consumption of sodium in grams/day (16).

Intake values above the Tolerable Upper Intake Level (UL) for age (17) were considered inadequate. The cut-off for the UL was based on the high sodium intake of Brazilians (18, 19).

The typical physical activity of adolescents was assessed using the short and validated version of the Brazilian adolescents of the International Physical Activity Questionnaire (IPAQ), which classifies populations into three categories: insufficiently active, sufficiently active and very active (20). The survey included questions regarding the practice of sports and the daily hours of exposure to television, videogames and computers.

Asthma characteristics were evaluated by the *International study of asthma and allergies in childhood* (ISAAC) self-administered questionnaire, which was validated after its translation into Brazilian Portuguese (21).

Asthma was assessed by the overall ISAAC score, as recommended by Wandalsen et al. (22). To identify adolescents with asthma, we used a cut-off score ≥ 6 on the ISAAC questionnaire (21). More severe asthma cases were defined by wheezing in the last 12 months and one or more positive responses to the following questions: inability to speak at least two complete words during an attack of wheezing, more than 12 crises in the last year, and greater than one night per week of nocturnal awakenings (23, 24).

Statistical analyses

The results are expressed as the mean and standard deviation. To assess the distribution of variables, the Kolmogorov-Smirnov test was used. To analyze the association between qualitative variables, the chi-square test was applied. Gross and adjusted estimation of prevalence ratios were calculated using a Poisson regression analysis with a confidence interval of 95% (95% CI) and a robust adjustment of variance in bivariate and multivariate analyses, respectively.

Covariates with a value of $p < 0.20$ in the bivariate analysis were used in the Poisson multiple regression analysis (multivariate analysis).

The cut-off level for significance was 5%, and the analyses were performed using SPSS (*Statistical Package for the Social Sciences*) software version 18.0.

RESULTS

We evaluated 1362 students from 57 classes across 16 high schools in the city of Santa Maria/RS. The average age of the adolescents was 15.65 ± 1.24 years, and the prevalence of signs and symptoms related to asthma among all adolescents were stratified by BMI-Z (Table 1). There were no significant differences in the signs and symptoms of asthma between adolescents with and without obesity.

Table 2 shows the physical activity and nutritional characteristics of the study participants. There is greater prevalence of obesity in females. There were statistically significant differences between overweight adolescents and adolescents with asthma, particularly in females. A total of 57.1% of the overweight adolescents with severe asthma consumed a high amount of sodium on a daily basis.

Based on the height-for-age index, we determined that out of 22 adolescents with short stature, 12 were classified as overweight. Among asthmatic students, there was no individual that was overweight and had a short stature.

The bivariate and multivariate analyses adjusted for the age of all adolescents, showed a significant risk

Table 1. The prevalence of signs and symptoms related to asthma in overweight and non-overweight adolescents according to the ISAAC questionnaire.

Questionnaire	Adolescents			
	Total* (n=1362)	Non-overweight* (n=995)	Overweight* (n=367)	P
Wheezing or whistling at any time	588 (43,2)	424(51,1)	164(12,0)	0,493
Wheezing or whistling in the chest	282 (20,7)	221(15,5)	71(5,2)	0,452
Attacks of wheezing	38 (2,8)	25(1,8)	71(5,2)	0,306
Sleep been disturbed due to wheezing	179 (13,1)	136(10,0)	43(3,2)	0,344
Wheezing to limit your speech	46 (3,4)	33(2,4)	13(1,0)	0,848
Ever asthma	219 (16,1)	149(10,9)	70(5,1)	0,069
Wheezy during or after exercise	217 (16)	162(12,0)	55(4,1)	0,564
Dry cough at night	491 (36,2)	360(26,5)	131(9,7)	0,882
Score and cut-off point (≥ 6)	279(20,5)	207(15,2)	72(5,3)	0,796
Severe asthma	78(5,7)	56(4,1)	22(1,6)	0,631

ISAAC, International study of asthma and allergies in childhood.*Number of subjects (percentage), n(%); Data were analyzed with a Pearson's chi-square test, ** $p < 0.05$.

Table 2. The effect of asthma on the nutritional status and physical activity level of adolescents.

Variables		Adolescents		Asthma		Severe asthma	
		Total (n=1362)	Overweight* (n=367)	Total (n=279)	Overweight* (n=72)	Total (n=78)	Overweight* (n=22)
Sex	Male	574(42.1)	155(42.2)	85(30.5)**	22(30.6)**	23(29.5)**	7(31.8)
	Female	788(57.9)	212(57.8)	194(69.5)	50(69.4)	55(70.5)	15(68.2)
H/A-Z	Appropriate	1340(98.4)	335(96.7)**	273(97.3)	72(100)	74(94.9)**	22(100)
	Short	22(1.6)	12(3.3)	6(2.2)	0	4(5.1)	0
WC	Without excess	862 (63.3)	224(61)	182(65.2)	44(61.1)	49(62.8)	12(54.5)
	With excess	500(36.7)	143(39)	97(34.8)	28(38.9)	29(37.2)	10(45.5)
AC	Eutrophic	792 (82.2)	212(82.8)	163(81.5)	41(77.4)	50(87.7)	14(87.5)
	Obesity	172(17.8)	44(17.2)	37(18.5)	12(22.6)	7(12.3)	2(12.5)
IPAC	Insuff. active	806 (67.4)	210(67.3)	172(65.6)	47(69.1)	46(63.9)	14(67.1)
	Suff. active	270 (22.6)	72(23.1)	61(23.3)	14(20.6)	20(27.8)	4(23.3)
	Very active	120 (10)	30(9.6)	29(24.2)	7(10.3)	6(8.3)	2 (9.6)
Tv/video	<4 hours	363(30.3)	84(26.8)	73(27.9)	17(25)	21(29.2)	3(15)
	>4 hours	834(69.7)	229(73.2)	189(72.1)	51(75)	51(70.8)	17(85)
Sports	No	545(45.7)	134(43.1)	127(48.5)	26(38.2)	34(47.2)	7(35)
	Yes	648(54.3)	177(56.9)	135(51.5)	42(61.8)	38(52.8)	13(65)
Sodium/day	Adequate	716(74.3)	191(75.5)	124(70.1)	25(54.3)**	28(57.1)**	6(42.9)**
	High	248(25.7)	78(24.5)	53(29.9)	21(45.7)	21(42.9)	8(57.1)

Index height-for-age (H/A-Z), Waist circumference (WC), Arm circumference (AC), International Physical Activity Questionnaire (IPAQ), *Number of subjects (percentage), n (%). Data were analyzed using a Pearson's chi-square test, **p<0.05.

for asthma in females and severe asthma in adolescents with high sodium intake (Table 3).

As demonstrated in Figure 1 and Table 4, overweight adolescents that were female and had a high sodium intake showed a significant risk of asthma in the

bivariate analysis. However, in the multivariate analysis, only high sodium intake was associated with asthma. The multivariate analysis also showed that severe asthma was more prevalent in adolescents with a high sodium intake.

Table 3. The risk factors for asthma and asthma severity among adolescents identified with bivariate and multivariate analyses

Variables	Asthma				Severe asthma					
	Total* (n=1362)		Total** (n=1362)		Total* (n=1362)		Total** (n=1362)			
	PR	CI 95%	P	PR	CI 95%	P	PR	CI 95%	P	
BMI-Z										
Non-overweight	Ref.	--	-	-	--	-	Ref.	--	-	-
Overweight	0.94	0.74-1.20	0.623			0.793	1.07	0.66-1.72	0.793	
Sex										
Male	Ref.	--	-	Ref.	--	-	Ref.	--	-	-
Female	1.66	1.32-2.09	<0.01 [†]	1.41	1.06-1.87	0.017 [†]	1.75	1.09-2.80	0.021 [†]	1.57
WC										
Without excess	Ref.	--	-	-	--	-	Ref.	--	-	-
With excess	0.92	0.74-1.15	0.473			0.938	1.01	0.65-1.58	0.938	
AC										
Eutrophic	Ref.	--	-	-	--	-	Ref.	--	-	-
Obesity	1.04	0.76-1.42	0.823			0.258	0.64	0.29-1.38	0.258	
TV/video										
<4 hours	Ref.	--	-	-	--	-	Ref.	--	-	-
>4 hours	1.12	0.88-1.43	0.341			0.832	1.05	0.64-1.73	0.832	
IPAC										
Insuff. active	Ref.	--	-	-	--	-	Ref.	--	-	-
Suff. active	0.93	0.63-1.37	0.734			0.385	1.48	0.61-3.59	0.385	
Very active	0.88	0.63-1.24	0.477			0.755	1.14	0.49-2.61	0.755	
Sports										
Yes	Ref.	--	-	-	--	-	Ref.	--	-	-
No	1.12	0.90-1.38	0.298			0.784	1.06	0.68-1.66	0.784	
Sodium/day										
Adequate	Ref.	--	-	Ref.	--	-	Ref.	--	-	-
High	1.24	0.93-1.65	0.146	1.29	0.96-1.75	0.083	2.16	1.24-3.74	0.006	2.30
										1.37-4.17
										0.006 [†]

Waist circumference (WC), Arm circumference (AC), International Physical Activity Questionnaire (IPAQ), 95% confidence interval (CI 95%), prevalence ratio (PR), reference (Ref.). Poisson regression: *bivariate analysis, ** multiple regression, [†]p<0,05.




Figure 1. The factors associated with asthma and asthma severity among overweight were identified with bivariate analysis.

Waist circumference (WC), Arm circumference (AC), International Physical Activity Questionnaire (IPAQ), 95% confidence interval (CI95%), prevalence ratio (PR). Poisson regression: *bivariate analysis, ** multiple regression, †p<0,05.

DISCUSSION

In this study, female sex and high sodium intake increased the risk for asthma. There was an additional risk for asthma in females and those with high sodium intake among overweight adolescents. There were no differences in the signs and symptoms of asthma between overweight and non-overweight adolescents; however, females with asthma showed a higher prevalence of obesity.

Asthma and obesity are among the most prevalent diseases in children. Both pathologies have increased in parallel over the recent decades, particularly in western countries. Therefore, the possibility of the association and risk between asthma and obesity has garnered a great deal of attention. Many studies used cross-sectional design to show an association between asthma and obesity in the adult population, especially in women. This association in

Table 4. The risk factors for asthma and asthma severity among overweight adolescents identified with a multivariate analysis

Variables	Asthma			Severe asthma		
	Overweight* (n=367)			Overweight* (n=367)		
	PR	CI95%	P	PR	CI95%	P
Sex						
Male	Ref.	--	-	-	--	-
Female	1.03	0.59-1.78	0.911			
AC						
Eutrophic	Ref.	--	-	-	--	-
Obesity	1.71	0.98-2.97	0.058			
TV/video						
<4 hours	-	--	-	Ref.	--	-
>4 hours				3.84	0.49-2.95	0.196
Sodium/day						
Adequate	Ref.	--	-	Ref.	-	-
High	2.48	1.46-4.19	0.001 [†]	3.07	1.03-9.13	0.043 [†]

Arm circumference (AC), 95% confidence interval (CI95%), prevalence ratio (PR), reference (Ref.).*adjusted for age, multiple regression-Poisson, [†]p<0,05.

children and adolescents remains contradictory (3, 21).

Recently, cross-sectional studies and studies with a prospective design (25-28, 31-33) reported an association between asthma and obesity in children and adolescents, especially among women (31, 33). Notably, the 95th percentile for BMI was used as the cut-off for nutritional status (25-27). These results are consistent with the hypothesis described in the meta-analyses completed by Beuther and Sutherland (36) and Chen et al. (37), which concludes that the incidence of asthma is weight-dependent. In contrast, other authors (29, 30, 34, 35) did not observe this association in the pediatric population. However, these authors used the 85th percentile for BMI as the cut-off point. This cut-off value corresponds to a z-score +1, which was the cut-off value used in the present study.

Our findings are in accordance with other studies (38, 39) that showed a higher prevalence of asthma or severe asthma in females (OR= 1.48 to 2.44). In the current study, the prevalence rate for asthma in all female participants and in all overweight adolescents was 1.66. This meta-analysis also showed that overweight/obese males are more susceptible to asthma (RR: 2.47 for boys, 1.25 for girls); however, a weight-dependence for the incidence of asthma was only significant in girls (37).

There is a significant debate regarding the effect of gender on asthma. Previous studies showed that boys have a higher incidence of asthma in childhood, but girls show the higher incidence during adolescence. Potential mechanisms to explain this phenomenon include pulmonary mechanics, sleep disorders and leptin, and the influence of hormones, such as estrogen (37, 40).

Notably, the majority of studies used BMI as the sole indicator of nutritional status. New findings indicate that body composition, including WC, in the pediatric population may increase the risk of asthma (3); however, we did not observe this characteristic in the present study. Interestingly, there was a tendency toward a positive risk for asthma based on the AC within overweight adolescents. This indicator reflects body composition, and it can be used as a practical and feasible measurement during clinical examination, especially when weight and height data are not available (41).

Possible explanations for the relationship between excess adipose tissue and asthma include genetic factors, mechanical changes, airway hyperresponsiveness, alteration of hormonal substances, such as cytokines and chemokines, and changes in lifestyle, including physical activity and nutrition (42).

Another indicator of nutritional status is the H/A index. A total of 2% of adolescents with asthma and 5% of adolescents with severe asthma also had a short stature; however, no adolescent with a small stature was overweight. These results are consistent with the study by Baum et al. (43) that reported children with allergy-related diseases, including asthma, are more likely to have a low height for their age. Asthmatics show a characteristic pattern of growth, *i.e.*, decreased growth rate, especially patients with severe asthma (44). Studies suggest that the use of inhaled corticosteroids at the recommended doses does not adversely affect the final growth of children and adolescents (45, 46).

The nutritional status of adolescents is likely related to asthma, and many theories have been proposed to elucidate the mechanism of this association. Dietary measures should be considered in analyses because the energetic control and quality of diet are considered possible inflammatory stimuli. Few studies have investigated the expenditure and consumption of energy, which are determinants of energy balance and

body weight. In children and adolescents with asthma, we found that the resting energy expenditure is similar between children with and without asthma; however, the estimated energy intake exceeds the resting energy expenditure in children with asthma (47).

These data reinforce that the increase in the prevalence of asthma, especially in westernized countries, is associated with environmental factors, such as dietary intake and sedentary lifestyle. Indeed, these elements likely play a fundamental role in the development and progression of asthma (48).

There was no statistically significant change in the risk of asthma based on the level of physical activities, sedentary activities and sports practice; however, the risk for severe asthma was higher among students classified as insufficiently active, and the risk doubled for overweight students that spent more than 4 hours a day doing sedentary activities. A previous study in children found no association between regular sport activity and asthma symptoms, but children who spent five or more hours per day watching television were more likely to have wheezing and asthma (48). Cabral et al. (49) studied children with asthma and found a higher prevalence of exercise-induced bronchospasm in more serious asthma cases; however, the intensity of the bronchoconstriction response to exercise was not related to asthma severity. Rodrigues et al. (50) observed that obese students showed more hyperresponsiveness to exercise. The authors did not consider this a reason to limit physical activity but rather suggested that this limitation be combined with other factors related to adipose tissue.

Notably, over 70% of the overweight adolescents evaluated in this study watched television or played videogames for more than four hours per day. The exposure to television is associated with an increase in the consumption of salty snacks, and these two behaviors are implicated in both obesity and the development of asthma (11) (5).

The consumption of fast foods that are high-calorie and rich in sodium and saturated/trans-fat is associated with the symptoms of wheezing and bronchial hyperresponsiveness in children; thus, diet plays an important role in the pathogenesis of asthma (5, 48).

The risk of asthma and increased asthma severity in all participants with high sodium intake was approximately two-fold higher. For overweight children, this risk was three-fold higher. Corroborating these findings, previous studies showed that the addition of salt to food led to a greater than two-fold increased risk for wheezing and asthma (48). The consumption of snacks (three times/week) was associated with asthma symptoms at a risk of 4.8 (5).

The influence of sodium on asthma is not yet fully understood. One potential explanation is that sodium exacerbates asthma via the hypersensitization of bronchial smooth muscle. This muscle is permeable to sodium, and an increase in sodium consumption leads to muscle hyperpolarization. Another hypothesis is that the asthma is associated with increased ouabain, an inhibitor of the sodium/potassium pump that causes sodium intracellular accumulation. However, the relationship between the response capacity of the airways (asthma) and urinary sodium excretion (an indicator of intake) in response to increased sodium intake is unclear (12). Based on the currently available evidence, we cannot conclude that reducing salt in the diet will help treat or manage asthma symptoms; however, sodium intake among asthmatics should abide by the recommendations available for the age-group of the patient (17).

During adolescence, many changes occur that influence the development of particular lifestyles, including nutrition and physical activity. We are the first to report a Brazilian study that specifically investigated the prevalence of asthma based on the sodium intake of adolescents; however, one limitation of the study is that

the effects of asthma protective nutrients as potential confounders were not assessed. Another limitation is the lack of uniform criteria for the definition of asthma across studies. The ISAAC questionnaire is widely used for this purpose because of its specificity; however, the symptoms obtained by this survey are subject to biases because the information is self-reported.

CONCLUSIONS

This study found no differences between adolescents with asthma and obesity; however, the prevalence of asthma was higher in overweight females. There was a higher risk for females and those with a high sodium intake for asthma and severe asthma, respectively. Among overweight adolescents, there was an additional risk for females and those with a high sodium intake. Asthma and obesity are multifactorial diseases with environmental factors that are significant predictors of quality of life. Thus, we suggest further research is needed to clarify the relationship between asthma and obesity and the influence of sodium on this association. Adolescents need to learn about these risk factors because school is an ideal location for health promotion and risk prevention.

REFERENCES

1. Asher MI, Keil U, Anderson HR, Beasley R, Crane J, et al. International Study of Asthma and Allergies in Childhood (ISAAC): rationale and methods. *The European respiratory journal*. 1995;8(3):483-91. Epub 1995/03/01.
2. Gibson GJ. Obesity, respiratory function and breathlessness. *Thorax*. 2000;55 Suppl 1:S41-4. Epub 2000/08/16.
3. MUSAAD SM, PATTERSON T, ERICKSEN M, LINDSEY M, DIETRICH K et al. Comparison of anthropometric measures of obesity in childhood allergic asthma: central obesity is most relevant. *The Journal of allergy and clinical immunology*. 2009;123(6):1321-7 e12. Epub 2009/05/15.

4. Romieu I, Mannino DM, Redd SC, McGeehin MA. Dietary intake, physical activity, body mass index, and childhood asthma in the Third National Health And Nutrition Survey (NHANES III). *Pediatric pulmonology*. 2004;38(1):31-42. Epub 2004/06/02.
5. Arvaniti F, Priftis KN, Papadimitriou A, Yiallourous P, Kapsokefalou M et al. Salty-snack eating, television or video-game viewing, and asthma symptoms among 10- to 12-year-old children: the PANACEA study. *Journal of the American Dietetic Association*. 2011;111(2):251-7. Epub 2011/01/29.
6. Kumanyika SK. Minisymposium on obesity: overview and some strategic considerations. *Annual review of public health*. 2001;22:293-308. Epub 2001/03/29.
7. Monteiro CA, Mondini L, de Souza AL, Popkin BM. The nutrition transition in Brazil. *European journal of clinical nutrition*. 1995;49(2):105-13. Epub 1995/02/01.
8. Lang DM, Butz AM, Duggan AK, Serwint JR. Physical activity in urban school-aged children with asthma. *Pediatrics*. 2004;113(4):e341-6. Epub 2004/04/03.
9. Tsai HJ, Tsai AC, Nriagu J, Ghosh D, Gong M et al. Associations of BMI, TV-watching time, and physical activity on respiratory symptoms and asthma in 5th grade schoolchildren in Taipei, Taiwan. *The Journal of asthma : official journal of the Association for the Care of Asthma*. 2007;44(5):397-401. Epub 2007/07/07.
10. Bar-on ME. The effects of television on child health: implications and recommendations. *Archives of disease in childhood*. 2000;83(4):289-92. Epub 2000/09/22.
11. Wiecha JL, Peterson KE, Ludwig DS, Kim J, Sobol A et al. When children eat what they watch: impact of television viewing on dietary intake in youth. *Archives of pediatrics & adolescent medicine*. 2006;160(4):436-42. Epub 2006/04/06.
12. Hirota SA, Janssen LJ. Sodium and asthma: something borrowed, something new? *American journal of physiology Lung cellular and molecular physiology*. 2007;293(6):L1369-73. Epub 2007/10/02.
13. WHO. Development of a WHO growth reference for school-aged children and adolescents. 2007.
14. Frisancho AR. Triceps skin fold and upper arm muscle size norms for assessment of nutritional status. *The American journal of clinical nutrition*. 1974;27(10):1052-8.
15. Taylor RW, Jones IE, Williams SM, Goulding A. Evaluation of waist circumference, waist-to-hip ratio, and the conicity index as screening tools for high trunk fat mass, as measured by dual-energy X-ray absorptiometry, in children aged 3-19 y. *The American journal of clinical nutrition*. 2000;72(2):490-5. Epub 2000/08/02.
16. Ferreira-Sae MC, Gallani MC, Nadruz W, Rodrigues RC, Franchini KG et al. Reliability and validity of a semi-quantitative FFQ for sodium intake in low-income and low-literacy Brazilian hypertensive subjects. *Public health nutrition*. 2009;12(11):2168-73. Epub 2009/05/30.
17. IOM. Food and Nutrition Board. Dietary reference intakes for water, potassium, sodium, chloride, and sulfate. Washington (DC): National Academy Press; 2000.
18. Sarno F, Claro RM, Levy RB, Bandoni DH, Ferreira SRG et al. Estimativa de consumo de sódio pela população brasileira, 2002-2003. *Revista de Saúde Pública*. 2009;43:219-25.
19. Veiga GVd, Costa RSd, Araújo MC, Souza AdM, Bezerra IN et al. Inadequação do consumo de nutrientes entre adolescentes brasileiros. *Revista de Saúde Pública*. 2013;47:212s-21s.
20. Guedes DP, Lopes CC, Guedes JERP. Reprodução

- bilidade e validade do Questionário Internacional de Atividade Física em adolescentes. *Revista Brasileira de Medicina do Esporte*. 2005;11:151-8.
21. Sole D, Vanna AT, Yamada E, Rizzo MC, Naspitz CK. International Study of Asthma and Allergies in Childhood (ISAAC) written questionnaire: validation of the asthma component among Brazilian children. *Journal of investigational allergology & clinical immunology*. 1998;8(6):376-82. Epub 1999/02/24.
22. Wandalsen NF, Gonzalez C, Wandalsen GF, Solé D. Avaliação de critérios para o diagnóstico de asma através de um questionário epidemiológico. *Jornal Brasileiro de Pneumologia*. 2009;35:199-205.
23. Mallol J, Crane J, von Mutius E, Odhiambo J, Keil U et al. The International Study of Asthma and Allergies in Childhood (ISAAC) Phase Three: a global synthesis. *Allergologia et immunopathologia*. 2013;41(2):73-85. Epub 2012/07/10.
24. Simões SdM, Cunha SSd, Barreto ML, Cruz ÁA. Distribuição da gravidade da asma na infância. *Jornal de Pediatria*. 2010;86:417-23.
25. Visness CM, London SJ, Zeldin DC. Association of Childhood Obesity with Atopic and Non-Atopic Asthma: Results from the National Health and Nutrition Examination Survey 1999-2006. *The Journal of Asthma*. 2010;47(7):822-9.
26. Okabe Y, Itazawa T, Adachi Y, Yoshida K, Ohya Y et al. Association of overweight with asthma symptoms in Japanese school children. *Pediatrics International*. 2011;53(2):192-8.
27. Chu Y-T, Chen W-Y, Wang T-N, Tseng H-I, Wu J-R et al. Extreme BMI predicts higher asthma prevalence and is associated with lung function impairment in school-aged children. *Pediatric pulmonology*. 2009;44(5):472-9.
28. Cibella F, Cuttitta G, La Grutta S, Melis MR, Bucchieri S et al. A cross-sectional study assessing the relationship between BMI, asthma, atopy, and eNO among schoolchildren. *Annals of allergy, asthma & immunology : official publication of the American College of Allergy, Asthma, & Immunology*. 2011;107(4):330-6.
29. Leung TF, Kong AP, Chan IH, Choi KC, Ho CS et al. Association between obesity and atopy in Chinese schoolchildren. *International archives of allergy and immunology*. 2009;149(2):133-40. Epub 2009/01/08.
30. Bertolace MdPC, Toledo E, Jorge PPdO, Liberatore Junior RDR. Association between obesity and asthma among teenagers. *Sao Paulo Medical Journal*. 2008;126:285-7.
31. Kattan M, Kumar R, Bloomberg GR, Mitchell HE, Calatroni A et al. Asthma control, adiposity, and adipokines among inner-city adolescents. *The Journal of allergy and clinical immunology*. 2010;125(3):584-92. Epub 2010/03/17.
32. Ginde AA, Santillan AA, Clark S, Camargo CA, Jr. Body mass index and acute asthma severity among children presenting to the emergency department. *Pediatric allergy and immunology : official publication of the European Society of Pediatric Allergy and Immunology*. 2010;21(3):480-8. Epub 2009/06/25.
33. Lang JE, Hossain J, Dixon AE, Shade D, Wise RA, Peters SP, et al. Does age impact the obese asthma phenotype? Longitudinal asthma control, airway function, and airflow perception among mild persistent asthmatics. *Chest*. 2011;140(6):1524-33. Epub 2011/07/30.
34. Hom J, Morley EJ, Sasso P, Sinert R. Body mass index and pediatric asthma outcomes. *Pediatric emergency care*. 2009;25(9):569-71. Epub 2009/09/17.
35. Peters JI, McKinney JM, Smith B, Wood P, Forkner E

- et al. Impact of obesity in asthma: evidence from a large prospective disease management study. *Annals of allergy, asthma & immunology : official publication of the American College of Allergy, Asthma, & Immunology*. 2011;106(1):30-5. Epub 2011/01/05.
36. Beuther DA, Sutherland ER. Overweight, obesity, and incident asthma: a meta-analysis of prospective epidemiologic studies. *American journal of respiratory and critical care medicine*. 2007;175(7):661-6. Epub 2007/01/20.
37. Chen YC, Dong GH, Lin KC, Lee YL. Gender difference of childhood overweight and obesity in predicting the risk of incident asthma: a systematic review and meta-analysis. *Obesity reviews : an official journal of the International Association for the Study of Obesity*. 2013;14(3):222-31. Epub 2012/11/14.
38. Moreau D, Kalaboka S, Choquet M, Annesi-Maesano I. Asthma, obesity, and eating behaviors according to the diagnostic and statistical manual of mental disorders IV in a large population-based sample of adolescents. *The American journal of clinical nutrition*. 2009;89(5):1292-8. Epub 2009/03/27.
39. He QQ, Wong TW, Du L, Jiang ZQ, Qiu H et al. Respiratory health in overweight and obese Chinese children. *Pediatric pulmonology*. 2009;44(10):997-1002. Epub 2009/09/22.
40. Hancox RJ, Milne BJ, Poulton R, Taylor DR, Greene JM et al. Sex differences in the relation between body mass index and asthma and atopy in a birth cohort. *American journal of respiratory and critical care medicine*. 2005;171(5):440-5. Epub 2004/11/24.
41. Zemel BS, Riley EM, Stallings VA. Evaluation of methodology for nutritional assessment in children: anthropometry, body composition, and energy expenditure. *Annual review of nutrition*. 1997;17:211-35. Epub 1997/01/01.
42. Papoutsakis C, Priftis KN, Drakouli M, Prifti S, Konstantaki E et al. Childhood overweight/obesity and asthma: is there a link? A systematic review of recent epidemiologic evidence. *Journal of the Academy of Nutrition and Dietetics*. 2013;113(1):77-105. Epub 2012/12/25.
43. Baum WF, Schneyer U, Lantzsch AM, Kloditz E. Delay of growth and development in children with bronchial asthma, atopic dermatitis and allergic rhinitis. *Experimental and clinical endocrinology & diabetes : official journal, German Society of Endocrinology [and] German Diabetes Association*. 2002;110(2):53-9. Epub 2002/04/03.
44. Sant'Anna CA, Solé D, Naspitz CK. Short stature in children with respiratory allergy. *Pediatric Allergy and Immunology*. 2006;7(4):187-92.
45. Arend EE, Fischer GB, Mocelin H, Medeiros L. Corticóide inalatório: efeitos no crescimento e na supressão adrenal. *Jornal Brasileiro de Pneumologia*. 2005;31:341-9.
46. Bensch GW, Greos LS, Gawchik S, Kpamegan E, Newman KB. Linear growth and bone maturation are unaffected by 1 year of therapy with inhaled flunisolide hydrofluoroalkane in prepubescent children with mild persistent asthma: a randomized, double-blind, placebo-controlled trial. *Annals of allergy, asthma & immunology : official publication of the American College of Allergy, Asthma, & Immunology*. 2011;107(4):323-9. Epub 2011/10/04.
47. Benedetti FJ, Mocelin HT, Bosa VL, de Mello ED, Fischer GB. Energy expenditure and estimated caloric intake in asthmatic adolescents with excess body weight. *Nutrition (Burbank, Los Angeles County, Calif)*. 2010;26(10):952-7.
48. Corbo GM, Forastiere F, De Sario M, Brunetti L, Bonci E et al. Wheeze and asthma in children:

associations with body mass index, sports, television viewing, and diet. *Epidemiology*. 2008;19(5):747-55. Epub 2008/05/23.

49. Cabral AL, Conceicao GM, Fonseca-Guedes CH, Martins MA. Exercise-induced bronchospasm in children: effects of asthma severity. *American journal of respiratory and critical care medicine*. 1999;159(6):1819-23. Epub 1999/06/03.
50. Rodrigues JC, Takahashi A, Olmos FMA, Souza JBd, Bussamra MHF et al. Efeito do índice de massa corpórea na gravidade da asma e na reatividade brônquica induzida pelo exercício em crianças asmáticas com sobrepeso e obesas. *Revista Paulista de Pediatria*. 2007;25:207-13.